



SUNSPOT RESULTS FOR JANUARY 2017

All observations carried out by HOWARD BARNES .

Telescope : 76 mm refractor (f.l. 910 mm) k considered as 1 .

Observed by PROJECTION . Full disc diameter = 145 mm approx .

WN = Wolf Number ; PX = Pettisindex ; BX = Beckindex ; CV = Classification Value ;

QC = Quality Count ; QC² = Squared Quality Count .

Q = Quietness [ie. steadiness] refer to Kiepenheuer scale .

S = Sharpness [ie. clarity] refer to Kiepenheuer scale .

T = Transparency where 1 = excellent , 5 = worthless .

PLEASE NOTE:
 From July 2015's issue,
 the observed Pettisindex
 is labelled PX.

Stated times (UT) approximate Co-ordinated Universal Time / Temps Universel Coordonné (UTC).

DATE	UT	g	f	WN	p	s	PX	BX	CV	QC	QC ²	Q	S	T	Ref.											
01																										
02																										
03																										
04	1940	0	0	0	0	0	0	0	0	0	0	1.5	2.5	2.0	5938-5											
05	1935	0	0	0	0	0	0	0	0	0	0	1.0	2.5	2.5	5939-5											
06																										
07																										
08																										
09																										
10																										
11	1940	0	0	0	0	0	0	0	0	0	0	1.5	2.0	2.0	5940-6											
12																										
13																										
14																										
15																										
16	1945	2	5	25	3	1	31	109	32	6	20	1.5	3.0	3.0	5941-6											
17																										
18																										
19	2015	2	6	26	3	2	32	88	33	7	25	1.5	2.5	2.0	5942-6											
20																										
21																										
22																										
23	2040	2	9	29	4	2	42	354	45	8	32	2.0	2.5	2.0	5943-6											
24																										
25	2020	3	14	44	4	8	48	384	44	10	36	2.0	3.0	2.5	5944-6											
26																										
27																										
28	2010	2	7	27	3	3	33	145	32	6	20	1.5	3.0	2.5	5945-6											
29	2015	2	5	25	2	2	22	42	19	5	13	1.5	2.5	2.5	5946-6											
30																										
31																										
Σ	—	13	46	176	19	18	208	1122	205	42	146	14.0	23.5	21.0	—											
NOBS	—	9	9	9	9	9	9	9	9	9	9	9	9	9	—											
MNS	—	1.44	5.11	19.56	2.11	2.00	23.11	124.67	22.78	4.67	16.22	1.56	2.61	2.33	—											
MEAN WEIGHT =		0.4663					MEAN CONDITION =					2.1667					TRUNCATED WOLF NUMBER =					18.22				

Georgi Dobrovolski Solar Observatory



SUNSPOT DISTRIBUTION & INTER-SOL INDICES FOR JANUARY 2017

All observations carried out by HOWARD BARNES .
 Telescope : 76 mm refractor (f. l. 910 mm) .
 Observed by PROJECTION . Full disc diameter = 145 mm approx .
 IS = Inter-Sol Index .
 gr = number of multi-spot groups .
 grfp = number of umbrae within penumbrae within the groups (gr) .
 grf = number of non-penumbral spots within the groups (gr) .
 efp = number of single penumbral spots .
 ef = number of single non-penumbral spots .
 Q = Quietness [ie. steadiness] refer to Kiepenheuer scale .
 S = Sharpness [ie. clarity] refer to Kiepenheuer scale .
 T = Transparency where 1 = excellent , 5 = worthless .

DATE	UT	IS	gr	grfp	grf	efp	ef	Q	S	T	Ref.
01											
02											
03											
04	1940	0	0	0	0	0	0	1.5	2.5	2.0	5938-5
05	1935	0	0	0	0	0	0	1.0	2.5	2.5	5939-5
06											
07											
08											
09											
10											
11	1940	0	0	0	0	0	0	1.5	2.0	2.0	5940-6
12											
13											
14											
15											
16	1945	6	1	3	1	1	0	1.5	3.0	3.0	5941-6
17											
18											
19	2015	8	2	4	2	0	0	1.5	2.5	2.0	5942-6
20											
21											
22											
23	2040	11	2	7	2	0	0	2.0	2.5	2.0	5943-6
24											
25	2020	17	3	6	8	0	0	2.0	3.0	2.5	5944-6
26											
27											
28	2010	8	1	3	3	1	0	1.5	3.0	2.5	5945-6
29	2015	6	1	2	2	1	0	1.5	2.5	2.5	5946-6
30											
31											
Σ	—	56	10	25	18	3	0	14.0	23.5	21.0	—
NOBS	—	9	9	9	9	9	9	9	9	9	—
MNS	—	6.22	1.11	2.78	2.00	0.33	0.00	1.56	2.61	2.33	—

Georgi Dobrovolski Solar Observatory



SUNSPOT CENSUS BY CLASSIFICATION FOR JANUARY 2017

All observations carried out by HOWARD BARNES .
 Telescope : 76 mm refractor (f.l. 910 mm).
 Observed by PROJECTION . Full disc diameter = 145 mm approx .
 IF 2 OR MORE REGIONS ARE OF THE SAME CLASSIFICATION , THEN SUNSPOT COUNTS
 ARE SEPARATED BY SOLIDI (/) .

DATE	UT	A		B		C		D		E		F		G		H		J	
		g	f	g	f	g	f	g	f	g	f	g	f	g	f	g	f	g	f
01																			
02																			
03																			
04	1940	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
05	1935	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
06																			
07																			
08																			
09																			
10																			
11	1940	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
12																			
13																			
14																			
15																			
16	1945	0	0	0	0	0	0	1	4	0	0	0	0	0	0	0	0	1	1
17																			
18																			
19	2015	0	0	0	0	1	2	1	4	0	0	0	0	0	0	0	0	0	0
20																			
21																			
22																			
23	2040	0	0	0	0	0	0	1	3	0	0	0	0	1	6	0	0	0	0
24																			
25	2020	0	0	1	2	0	0	1	7	0	0	0	0	1	5	0	0	0	0
26																			
27																			
28	2010	0	0	0	0	0	0	1	6	0	0	0	0	0	0	0	0	1	1
29	2015	0	0	0	0	1	4	0	0	0	0	0	0	0	0	0	0	1	1
30																			
31																			
TOTALS	—	0	0	1	2	2	6	5	24	0	0	0	0	2	11	0	0	3	3
REGIONAL PERCENTAGES																			
A	B	C	D	E	F	G	H	J	Σg										
0.0	7.7	15.4	38.5	0.0	0.0	15.4	0.0	23.1	13										
NOBS = 9				$\overline{p/g}$ mean = 1.4722						$\overline{f/g}$ mean = 3.4444									
				$\overline{p/g}$ mean = 1.4615						$\overline{f/g}$ mean = 3.5385									
GROUP COMPLEXITY INDEX (GCI) = 5.0000																			



SMOOTHED RESULTS OF OBSERVED VALUES FOR THE LAST 12 MONTHS (OBTAINABLE) USING THE WALDMEIER & BARNES-13 METHODS.

DATA BELOW ARE PRELIMINARY. FINAL VALUES WILL BE PUBLISHED IN GDSO ANNUAL REPORTS.

WALDMEIER METHOD

MONTH	$g^r(S^w)$	$WN^r(S^w)$	$PX^r(S^w)$	$BX^r(S^w)$	$CV^r(S^w)$	$QC^r(S^w)$	$IS^r(S^w)$
2015 AUGUST	2.70	43.85	53.60	446.5	54.36	8.86	18.72
SEPTEMBER	2.72	43.83	53.41	443.9	53.62	8.89	18.50
OCTOBER	2.69	42.82	51.91	427.2	52.19	8.75	17.79
NOVEMBER	2.61	41.52	50.26	407.9	51.92	8.57	17.23
DECEMBER	2.51	39.57	47.78	374.0	50.79	8.28	16.24
2016 JANUARY	2.30	36.14	43.49	334.6	46.88	7.58	14.69
FEBRUARY	2.22	34.30	40.69	297.2	44.01	7.23	13.60
MARCH	2.22	33.19	38.55	259.5	42.02	7.04	12.52
APRIL	2.17	32.24	37.29	247.2	40.89	6.80	12.02
MAY	2.13	31.38	36.01	234.1	39.81	6.59	11.56
JUNE	2.05	29.80	33.32	204.2	37.05	6.23	10.70
JULY	1.97	28.48	31.41	185.6	34.55	5.94	10.11

BARNES-13 METHOD

MONTH	$g^r(S^{B13})$	$WN^r(S^{B13})$	$PX^r(S^{B13})$	$BX^r(S^{B13})$	$CV^r(S^{B13})$	$QC^r(S^{B13})$	$IS^r(S^{B13})$
2015 AUGUST	2.82	46.86	57.83	509.5	57.90	9.41	20.60
SEPTEMBER	2.72	45.08	55.86	490.1	55.87	9.15	19.78
OCTOBER	2.60	42.49	52.59	451.2	52.93	8.74	18.37
NOVEMBER	2.47	39.93	49.32	408.4	50.56	8.33	16.95
DECEMBER	2.38	37.61	46.09	360.9	48.30	7.94	15.49
2016 JANUARY	2.28	35.15	42.32	310.6	45.29	7.48	13.92
FEBRUARY	2.23	33.42	39.24	266.3	42.84	7.14	12.63
MARCH	2.21	32.39	37.16	233.9	41.44	6.93	11.74
APRIL	2.19	31.84	36.06	220.0	40.87	6.77	11.37
MAY	2.16	31.26	34.98	210.5	39.99	6.59	11.11
JUNE	2.09	30.22	33.33	199.1	38.04	6.31	10.71
JULY	2.02	29.31	32.23	195.0	36.19	6.07	10.48

NB: VALUES FROM MAY 2015 TO MAY 2016 (INCLUSIVE),
ARE BASED, IN PART, ON INTERPOLATED VALUES OF NOVEMBER 2015.